

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

BEST AVAILABLE COPY**Amendments to the Claims:**

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (currently amended) A method of detecting or identifying an analyte of interest in a sample, comprising:
 - (i) contacting the sample containing the analyte with one or more affinity molecules to form a complex of the analyte and the one or more affinity molecules, wherein each of the one or more affinity molecules has an affinity against the analyte;
 - (ii) separating the complex and any unbound affinity molecule ~~in the presence of a charged polymer by using a separation channel in a microfluidic device, the separation channel being filled with a separation media and a charged polymer, the microfluidic device comprising~~ at least one separation channel having at least one microscale dimension of between about 0.1 and 500 microns; and
 - (iii) detecting the complex to identify the presence of the analyte or to determine an amount of the analyte in the sample, wherein the charged polymer reduces interference with detecting.
2. (original) The method of claim 1, wherein the charged polymer is a polyanionic polymer or a polycationic polymer.
3. (original) The method of claim 2, wherein the charged polymer is a polyanionic polymer selected from the group consisting of polysaccharides, polynucleotides, polypeptides, synthetic macromolecular compounds, ceramics and a complex thereof.
4. (original) The method of claim 1, wherein the charged polymer is a polyanionic polymer selected from the group consisting of poly-dIdC, heparin sulfate, dextran sulfate, polytungstic acid, polyanethole sulfonic acid, polyvinyl sulfate, polyacrylate, chondroitin sulfate, plasmid DNA, calf thymus DNA, salmon sperm DNA, DNA coupled to cellulose, glass particles, colloidal glass, and glass milk.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

5. (withdrawn) The method of claim 1, wherein the charged polymer is a polycationic polymer selected from the group consisting of polyallylamines, polylysine, polyhistidine, chitosan, protamine, polyethyleneimine and polyarginine.

6. (original) The method of claim 1, wherein the charged polymer comprises a net negative charge.

7. (withdrawn) The method of claim 1, wherein the charged polymer comprises a net positive charge.

8. (original) The method of claim 6, wherein the charged polymer comprises heparin sulfate.

9. (original) The method of claim 1, wherein at least one of the one or more affinity molecules is labeled with a detectable marker.

10. (original) The method of claim 1, wherein at least one of the one or more affinity molecules is bound to a charged carrier molecule to form a conjugate of the affinity molecule and the charged carrier molecule, and wherein the charged carrier molecule causes a change in a separation property of the analyte by binding to the analyte through the one or more affinity molecule to form a complex of the analyte, the affinity molecule and the charged carrier molecule.

11. (original) The method of claim 1 or 10, wherein the affinity molecule is one which binds to the analyte by a protein-protein interaction, a protein-chemical interaction or a chemical-chemical interaction.

12. (original) The method of claim 1 or 10, wherein the affinity molecule is one which binds to the analyte by an antigen-antibody interaction, a sugar chain-lectin interaction, an enzyme-inhibitor interaction, a protein-peptide chain interaction, a chromosome or nucleotide chain-nucleotide chain interaction, a nucleotide-ligand interaction or a receptor-ligand interaction.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

13. (original) The method of claim 1 or 10, wherein the affinity molecule is selected from the group consisting of an antibody, an Fab, F(ab')₂ or Fab' fragment of an antibody, an antibody variable region, a lection, avidin, a receptor, an affinity peptide, an aptamer, and a DNA binding protein.

14. (original) The method of claim 10, wherein the charged carrier molecule is an anionic molecule or a cationic molecule.

15. (original) The method of claim 14, wherein the charged carrier molecule is one having the same net charge as the charged polymer.

16. (original) The method of claim 14, wherein the charged carrier molecule is an anionic molecule comprising a nucleotide chain or a sulfonated polypeptide.

17. (original) The method of claim 10, wherein the charged carrier molecule comprises DNA, RNA, a cationic polymer, or a sulfonated polypeptide

18. (original) The method of claim 17, wherein the charged carrier molecule comprises DNA comprising one or more synthetic sequences.

19. (original) The method of claim 18, wherein the one or more synthetic sequences comprise one or more nucleotide analogs comprising a linker group or a linker reactive group.

20. (original) The method of claim 19, wherein the linker group or linker reactive group comprises an amino group, a thiol, a carboxyl group, an imidazol group, or a succinimide group.

21. (original) The method of claim 20, further comprising covalently bonding a detectable marker to the linker group or linker reactive group.

22. (currently amended) The method of claim 1, wherein at least one of the one or more affinity molecules is labeled with a detectable marker.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

23. (original) The method of claim 10, wherein at least one conjugate or at least one affinity molecule which does not form a conjugate is labeled with a detectable marker.

24. (original) The method of claim 10, wherein at least one affinity molecule and the charged carrier molecule forming the conjugate is labeled by a detectable marker.

25. (original) The method of claim 10, wherein the charged carrier molecule in the conjugate is labeled by a detectable marker.

26. (original) The method of claim 10, wherein the affinity molecule in the conjugate is labeled by a detectable marker.

27. (original) The method of claim 9, 21, 22, 23, 24, 25 or 26, wherein the detectable marker is a fluorescent dye, a luminescent dye, a phosphorescent dye, a fluorescent protein, a luminescent protein or particle, a radioactive tracer, a chemiluminescent compound, a redox mediator, an electrogenic compound, an enzyme, a colloidal gold particle, or a silver particle.

28. (original) The method of claim 10, wherein separating comprises electrophoretic separation of the conjugate or the complex through a separation media in the separation channel.

29. (original) The method of claim 28, wherein the separation media comprises a size exclusion resin, a polyacrylamide gel, polyethylene glycol (PEG), polyethyleneoxide (PEO), a co-polymer of sucrose and epichlorohydrin, polyvinylpyrrolidone (PVP), hydroxyethylcellulose (HEC), poly-N,N-dimethylacrylamide (pDMA), or an agarose gel.

30. (canceled)

31. (currently amended) The method of claim ~~30~~ 1, wherein the charged polymer is present in the separation media at a concentration of between about 0.01 to 5%.

32. (currently amended) The method of claim ~~30~~ 1, wherein the charged polymer is present in the separation media at a concentration of between about 0.05 to 2%.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

33. (original) The method of claim 28, further comprising introducing a charged polymer into a buffer which comprises the sample.

34. (original) The method of claim 33, wherein the charged polymer is present in the sample buffer at a concentration of between about 0.001 to 2%.

35. (original) The method of claim 1, wherein the separation channel has at least one cross-sectional microscale dimension of between about 0.1 and 200 microns.

36. (currently amended) The method of claim 1, wherein:

step (i) comprises contacting the sample containing the analyte with the one or more affinity molecules, at least one of ~~which is~~ the one or more affinity molecules being labeled by a detectable marker, to form a complex containing the analyte and the one or more affinity molecules labeled by the detectable marker;

step (ii) comprises separating the complex from any free affinity molecule labeled by the detectable marker ~~which that~~ is not involved in forming the complex in a the separation channel of a the microfluidic device in the presence of a charged polymer,

step (iii) comprises:

(a) measuring an amount of the separated complex or detecting a presence of the separated complex; and

(b) determining an amount of the analyte in the sample on the basis of the measured amount or identifying a presence of the analyte in the sample on the basis of the detected presence; and wherein each of the one or more affinity molecules has a property capable of binding to the analyte, ~~and wherein when two or more affinity molecules are used, each affinity molecule has a property capable of binding with the analyte at a different site on the analyte from every other affinity molecule.~~

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

37. (currently amended) The method of claim 10, wherein:

step (i) comprises contacting the sample containing the analyte with the one or more conjugates of ~~an~~the affinity molecule and ~~a~~the charged carrier molecule, wherein at least one of the one or more conjugates is labeled by a detectable marker, to form a complex containing the analyte and the conjugate labeled by the detectable marker;

step (ii) comprises separating the complex from the at least one conjugate labeled by the detectable marker ~~which~~that is not involved in forming the complex ~~in~~a the separation channel of ~~a~~the microfluidic device ~~in the presence of a charged polymer~~;

step (iii) comprises:

(a) measuring an amount of the separated complex or detecting a presence of the separated complex; and

(b) determining an amount of the analyte in the sample on the basis of the measured amount or identifying a presence of the analyte in the sample on the basis of the detected presence; and wherein the affinity molecule in the conjugate has a property capable of binding to the analyte, ~~and when two or more conjugates are used, each affinity molecule in the conjugate has a property capable of binding with the analyte at a different site on the analyte from every other affinity molecule, and the charged carrier molecule has a property capable of causing a change in a separation property of the analyte by binding to the analyte through the affinity molecule to form a complex of the analyte, the affinity molecule and the charged carrier molecule.~~

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

38. (currently amended) The method of claim 10, wherein:

step (i) comprises contacting the sample containing the analyte with the one or more affinity molecules and the one or more conjugates of ~~an~~the affinity molecule and ~~a~~the charged carrier molecule, wherein either at least one of the affinity molecules or at least one of the conjugates is labeled by a detectable marker, to form a complex containing the analyte, the affinity molecule, and the conjugate;

step (ii) comprises separating the complex from any free affinity molecule labeled by the detectable marker or the conjugate labeled by the detectable marker ~~which that~~ is not involved in forming the complex in ~~a~~the separation channel of ~~a~~the microfluidic device ~~in the presence of a charged polymer~~;

step (iii) comprises:

(a) measuring an amount of the separated complex or detecting a presence of the separated complex; and

(b) determining an amount of the analyte in the sample on the basis of the measured amount or identifying a presence of the analyte in the sample on the basis of the detected presence; and wherein each of the affinity molecule and the affinity molecule in the conjugate ~~have~~has a property capable of binding to the analyte, ~~and each affinity molecule has a property capable of binding with the analyte at a different site on the analyte from every other affinity molecule, and the charged carrier molecule has a property capable of causing a change in a separation property of the analyte by binding to the analyte through the affinity molecule to form a complex of the analyte, the affinity molecule and the charged carrier molecule.~~

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

39. (currently amended) A method for determining an analyte in a sample, ~~which comprises the method comprising:~~

(i) contacting the sample containing the analyte ~~with, either~~ the analyte labeled by a detectable marker or an analogue of the analyte labeled by a detectable marker, ~~and one or more~~ an affinity molecule to, thereby forming a first complex of the analyte in the sample and the affinity molecule and a second complex of either the labeled analyte and the affinity molecule or the labeled analogue and the affinity molecule;

(ii) separating the second complex from any free labeled analyte or the free labeled analogue ~~which that is not involved in forming the second complex in a separation channel of a microfluidic device in the presence of a charged polymer, the separation channel being filled with a separation media and a charged polymer;~~

(iii) measuring an amount of the separated second complex or an amount of the separated free labeled analyte or the separated free labeled analogue; and

(iv) determining an amount of the analyte in the sample on the basis of the measured amount; wherein the affinity molecule has a property capable of binding to the analyte in the sample and the labeled analyte or a property capable of binding to the analyte in the sample and the labeled analogue, ~~and wherein when two or more affinity molecules are used, each affinity molecule has a property capable of binding with the analyte in the sample and the labeled analyte at a different site on the analyte in the sample and a different site on the labeled analyte from every other affinity molecule or each affinity molecule has a property capable of binding with the analyte in the sample and the labeled analogue at a different site on the analyte in the sample and a different site on the labeled analogue from every other affinity molecule.~~

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

40. (currently amended) The method of claim 39, wherein:

step (i) comprises contacting the sample containing the analyte with either the labeled analyte labeled by a detectable marker or an the labeled analogue, of the analyte labeled by a detectable marker and one or more conjugate of an the affinity molecule and a charged carrier molecule ~~to, thereby forming a first complex of the analyte in the sample and the conjugate and a second complex of either the labeled analyte and the conjugate or the labeled analogue and the conjugate;~~

step (ii) comprises separating the second complex from any free labeled analyte or free labeled analogue ~~which that is not involved in forming the second complex in a the separation channel of a the microfluidic device in the presence of a charged polymer;~~

step (iii) comprises measuring an amount of the separated second complex or an amount of the separated free labeled analyte or the separated free labeled analogue; and

step (iv) comprises determining an amount of the analyte in the sample on the basis of the measured amount; and wherein the affinity molecule in the conjugate has a property capable of binding to the analyte in the sample and the labeled analyte or the analyte in the sample and the labeled analogue, ~~and when two or more conjugates are used, each affinity molecule in the conjugate has a property capable of binding with the analyte in the sample and the labeled analyte at a different site on the analyte in the sample and a different site on the labeled analyte from every other affinity molecule or each affinity molecule in the conjugate has a property capable of binding with the analyte in the sample and the labeled analogue at a different site on the analyte in the sample and a different site on the labeled analogue from every other affinity molecule, and wherein the charged carrier molecule has a property capable of causing a change in a separation property of the labeled analyte or the labeled analogue by binding to the labeled analyte or the labeled analogue through the affinity molecule to form a complex of the labeled analyte or the labeled analogue, the affinity molecule and the charged carrier molecule.~~

41. (currently amended) The method of claim 39, wherein:

step (i) comprises contacting the sample containing the analyte with either the labeled analyte labeled by a detectable marker or an the labeled analogue, of the analyte labeled

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

~~by a detectable marker, one or more~~ the affinity molecule, and ~~one or more~~ a conjugate of ~~an~~ the affinity molecule and a charged carrier molecule ~~to, thereby forming~~ a first complex of the analyte in the sample, the affinity molecule, and the conjugate and a second complex of either the labeled analyte, the affinity molecule, and the conjugate or the labeled analogue, the affinity molecule, and the conjugate;

step (ii) comprises separating the second complex from any free labeled analyte or the labeled analogue ~~which that is not involved in forming the second complex in a~~ the separation channel of ~~a~~ the microfluidic device ~~in the presence of a charged polymer;~~

step (iii) comprises measuring an amount of the separated second complex or an amount of the separated free labeled analyte or the separated free labeled analogue;

step (iv) comprises determining an amount of the analyte in the sample on the basis of the measured amount; and wherein each of the affinity molecule and the affinity molecule in the conjugate ~~have~~ has a property capable of binding to the analyte in the sample and the labeled analyte or the analyte in the sample and the labeled analogue, ~~and each affinity molecule has a property capable of binding with the analyte in the sample and the labeled analyte at the different site on the analyte in the sample and a different site on the labeled analyte from every other affinity molecule or each affinity molecule has a property capable of binding with the analyte in the sample and the labeled analogue at a different site on each of the analyte in the sample and a different site on the labeled analogue from every other affinity molecule, and wherein the charged carrier molecule has a property capable of causing a change in a separation property of the labeled analyte or the labeled analogue by binding to the labeled analyte or the labeled analogue through the affinity molecule to form a complex of the labeled analyte or the labeled analogue, the affinity molecule and the charged carrier molecule.~~

42. (currently amended) A method for determining an analyte in a sample, ~~which comprises the method comprising:~~

(i) contacting the sample containing the analyte ~~with, either~~ either the analyte bound to a charged carrier molecule or an analogue of the analyte bound to a charged carrier molecule, and ~~one or more~~ an affinity molecule labeled by a detectable marker ~~to, thereby forming~~ a first complex of either the analyte bound to the charged carrier molecule and the labeled affinity

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

molecule or the analogue bound to a the charged carrier molecule and the labeled affinity molecule and a second complex of the analyte in the sample and the labeled affinity molecule;

(ii) separating the first complex from any second complex in a separation channel of a microfluidic device in the presence of a charged polymer, the separation channel being filled with a separation media and a charged polymer;

(iii) measuring an amount of the separated first complex or an amount of the second complex;

(iv) determining an amount of the analyte in the sample on the basis of the measured amount; and wherein the affinity molecule has a property capable of binding to the analyte in the sample and the analyte bound to the charged carrier molecule or the analyte in the sample and the analogue bound to the charged carrier molecule, ~~and wherein when two or more affinity molecules are used, each affinity molecule has a property capable of binding with the analyte in the sample and the analyte bound to the charged carrier molecule at a different site on the analyte in the sample and a different site on the analyte bound to the charged carrier molecule from every other affinity molecule or each affinity molecule has a property capable of binding with the analyte in the sample and the analogue bound to the charged carrier molecule at a different site on the analyte in the sample and a different site on the analogue bound to the charged carrier molecule from every other affinity molecule, and wherein the charged carrier molecule has a property capable of causing a change in a separation property of the first complex by binding to the analyte or the analogue to form a complex of the analyte, the affinity molecule and the charged carrier molecule.~~

43. (original) The method of claim 1, wherein the sample comprises a serum, a plasma, a whole blood, a tissue extract, a cell extract, a nuclear extract, a culture media, a microbial culture extract, members of a molecular library, a clinical sample, a sputum specimen, a stool specimen, a cerebral spinal fluid, a urine sample, a uro-genital swab, a throat swab, or an environmental sample.

44. (original) The method of claim 1, wherein the analyte comprises AFP, hCG, TSH, FSH, LH, interleukin, Fas ligand, CA19-9, CA125, PSA, HBsAg, anti-HTV antibody, or T4.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

45-50 (canceled)

51. (currently amended) A method of concentrating an analyte of interest in a sample, the method comprising:

(i) contacting the sample containing the analyte with one or more of a conjugate of an affinity molecule and a charged carrier molecule to form a complex of the analyte and the conjugate;

(ii) concentrating the complex by using a concentration channel in a microfluidic device, the concentration channel being filled with a concentration media and a charged polymer, the microfluidic device comprising at least one concentration channel having at least one microscale dimension of between about 0.1 and 500 microns; and wherein the charged carrier molecule causes a change in a migration property of the analyte by binding to the analyte through the affinity molecule to form a complex of the analyte, the affinity molecule and the charged carrier molecule.

52. (canceled)

53. (original) The method of claim 51, wherein contacting the sample containing the analyte with one or more conjugate of an affinity molecule and a charged carrier molecule to form a complex of the analyte and the conjugate is conducted in a microchannel fluidically connected to the concentration channel having at least one microscale dimension of between about 0.1 and 500 microns.

54. (canceled)

55. (currently amended) The method of claim 51, wherein concentrating the complex is conducted by utilizing the difference in an electrophoretic mobility between the complex and ~~the noise constituents~~ in the sample on the basis of charge of the charged carrier molecule.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

56. (currently amended) The method of claim 51, wherein concentrating the complex is conducted by utilizing the difference in an adsorption property between the complex and the noise constituents in the sample on the basis of charge of the charged carrier molecule.

57. (original) The method of claim 51, wherein concentrating the complex is conducted according to a concentration method selected from the group consisting of field amplification sample stacking (FASS), field amplification sample injection (FASI), isotachophoresis (ITP), isoelectric focusing (IEF) and solid phase extraction (SPE).

58. (original) The method of claim 51, wherein concentrating the complex is conducted according to a concentration method selected from the group consisting of field amplification sample stacking (FASS) and isotachophoresis (ITP).

59. (original) The method of claim 51, wherein the charged carrier molecule is an anionic molecule or a cationic molecule.

60. (original) The method of claim 59, wherein the charged carrier molecule is an anionic molecule comprising a nucleotide chain or a sulfonated polypeptide.

61. (original) The method of claim 51, wherein the charged carrier molecule comprises DNA, RNA, a cationic polymer, or a sulfonated polypeptide

62. (original) The method of claim 61, wherein charged carrier molecule comprises DNA comprising one or more synthetic sequences.

63. (original) The method of claim 62, wherein the one or more synthetic sequences comprise one or more nucleotide analogs comprising a linker group or linker reactive group.

64. (original) The method of claim 63, wherein the linker group or linker reactive group comprises an amino group, a thiol, a carboxyl group, an imidazol group, or a succinimide group.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

65. (original) The method of claim 64, further comprising covalently bonding a detectable marker to the linker group or the linker reactive group.

66. (original) The method of claim 62, wherein the one or more synthetic sequence consists of one selected from a phosphorothioate analog of nucleotide, a nucleotide that contains a methylene group in the place of the oxygen in the ribose ring, or a nucleotide which has a replacement of the 2'-sugar deoxy substituent with 2'-fluoro, 2'-O-methyl, 2'-O-alkoxy]- and 2'-O-allyl modification

67. (original) The method of claim 51, wherein the contacting step further comprises contacting the sample with one or more affinity molecule to form a complex of the analyte, the conjugate and the affinity molecule.

68. (original) The method of claim 51 or 67, wherein the affinity molecule is one which binds to the analyte by a protein-protein interaction, a protein-chemical interaction or a chemical-chemical interaction.

69. (original) The method of claim 51 or 67, wherein the affinity molecule is one which binds to the analyte by an antigen-antibody interaction, a sugar chain-lectin interaction, an enzyme-inhibitor interaction, a protein-peptide chain interaction, a chromosome or nucleotide chain-nucleotide chain interaction, a nucleotide-ligand interaction or a receptor-ligand interaction.

70. (original) The method of claim 51 or 67, wherein the affinity molecule is selected from the group consisting of an antibody, an Fab, F(ab')₂ or Fab' fragment of an antibody, an antibody variable region, a lection, avidin, a receptor, an affinity peptide, an aptamer, and a DNA binding protein.

71. (original) The method of claim 67, wherein at least one conjugate or at least one affinity molecule which does not form a conjugate is labeled with a detectable marker.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

72. (original) The method of claim 51, wherein at least one of the affinity molecule and the charged carrier molecule forming the conjugate is labeled by a detectable marker.

73. (original) The method of claim 51, wherein the charged carrier molecule in the conjugate is labeled by a detectable marker.

74. (original) The method of claim 51, wherein the affinity molecule in the conjugate is labeled by a detectable marker.

75. (original) The method of claim 67, 71, 72, 73 or 74, wherein the detectable marker is a fluorescent dye, a luminescent dye, a phosphorescent dye, a fluorescent protein, a luminescent protein or particle, a radioactive tracer, a chemiluminescent compound, a redox mediator, an electrogenic compound, an enzyme, a colloidal gold particle, or a silver particle.

76. (original) The method of claim 51 or 67, wherein the contacting step and/or the concentrating step is conducted in the presence of a charged polymer.

77. (original) The method of claim 76, wherein the charged polymer is a polyanionic polymer or a polycationic polymer.

78. (original) The method of claim 77, wherein the charged polymer is a polyanionic polymer selected from the group consisting of polysaccharides, polynucleotides, polypeptides, synthetic macromolecular compounds, ceramics and complexes thereof.

79. (original) The method of claim 76, wherein the charged polymer is a polyanion selected from the group consisting of poly-dIdC, heparin sulfate, dextran sulfate, polytungstic acid, polyanethole sulfonic acid, polyvinyl sulfate, polyacrylate, chondroitin sulfate, plasmid DNA, calf thymus DNA, salmon sperm DNA, DNA coupled to cellulose, glass particles, colloidal glass, and glass milk.

10/821,657 filed 04/08/2004.

Wada et al.

Reply to Office Action of 07/24/2006

80. (withdrawn) The method of claim 76, wherein the charged polymer is a polycation selected from the group consisting of polyallylamine, polylysine, polyhistidine, chitosan, protamine, polyethyleneimine and polyarginine.

81. (original) The method of claim 76, wherein the charged polymer comprises a net negative charge.

82. (withdrawn) The method of claim 76, wherein the charged polymer comprises a net positive charge.

83. (original) The method of claim 76, wherein the charged carrier molecule and the charged polymer are the same net charge.

84. (original) The method of claim 81, wherein the charged polymer comprises heparin sulfate.

85. (original) The method of claim 51 or 67, wherein the concentrating step comprises electrophoretic concentration of the conjugate or the complex through a concentration media in the concentration channel.

86. (original) The method of claim 85, wherein the concentration media comprises a size exclusion resin, a polyacrylamide gel, polyethylene glycol (PEG), polyethyleneoxide (PEO), a co-polymer of sucrose and epichlorohydrin, polyvinylpyrrolidone (PVP), hydroxyethylcellulose (HEC), poly-N,N-dimethylacrylamide (pDMA), or an agarose gel.

87. (original) The method of claim 85, wherein the concentration media further comprises a charged polymer.

88. (original) The method of claim 87, wherein the charged polymer is present in the concentration media at a concentration of between about 0.01 to 5%.

89. (original) The method of claim 87, wherein the charged polymer is present in the concentration media at a concentration of between about 0.05 to 2%.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

90. (original) The method of claim 85, further comprising introducing a charged polymer into a buffer which comprises the sample.

91. (original) The method of claim 90, wherein the charged polymer comprises heparin sulfate which is present in the sample buffer at a concentration of between about 0.001 to 2%.

92. (original) The method of claim 51, wherein the concentration channel has at least one cross-sectional microscale dimension of between about 0.1 and 200 microns.

93. (currently amended) A method of detecting or identifying an analyte of interest in a sample, the method comprising:

(i) contacting the sample containing the analyte with one or more a conjugates of an affinity molecule and a charged carrier molecule to form a complex of the analyte and the conjugate;

(ii) concentrating the complex by using a concentration channel in a microfluidic device comprising at least one concentration channel having at least one microscale dimension of between about 0.1 and 500 microns;

(iii) separating the complex and any unbound conjugate by using a separation channel in a microfluidic device comprising at least one separation channel, the separation channel being filled with a separation media and a charged polymer, the separation channel having at least one microscale dimension of between about 0.1 and 500 microns ~~and comprising a charged polymer in the separation channel~~;

(iv) detecting the complex to identify the presence of the analyte or to determine an amount of the analyte in the sample, wherein the contacting, concentrating and/or separating step is conducted in the presence of a charged polymer which reduces interference with detecting; and wherein the charged carrier molecule causes a change in a migration property of the analyte by binding to the analyte through the affinity molecule to form a complex of the analyte, the affinity molecule and the charged carrier molecule.

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

94. (new) The method of claim 36, wherein two or more affinity molecules are used, and wherein each affinity molecule has a property capable of binding with the analyte at a different site on the analyte from every other affinity molecule.

95. (new) The method of claim 37, wherein two or more conjugates are used, and wherein each affinity molecule in the two or more conjugates has a property capable of binding with the analyte at a different site on the analyte from every other affinity molecule, and wherein the charged carrier molecule has a property capable of causing a change in a separation property of the analyte by binding to the analyte through the affinity molecule to form a complex of the analyte, the affinity molecule, and the charged carrier molecule.

96. (new) The method of claim 38, wherein two or more affinity molecules are used, and wherein each affinity molecule has a property capable of binding with the analyte at a different site on the analyte from every other affinity molecule, and wherein the charged carrier molecule has a property capable of causing a change in a separation property of the analyte by binding to the analyte through the affinity molecule to form a complex of the analyte, the affinity molecule, and the charged carrier molecule.

97. (new) The method of claim 39, wherein two or more affinity molecules are used, and wherein each affinity molecule has a property capable of binding with the analyte in the sample and the labeled analyte at a different site on the analyte in the sample and a different site on the labeled analyte from every other affinity molecule, or each affinity molecule has a property capable of binding with the analyte in the sample and the labeled analogue at a different site on the analyte in the sample and a different site on the labeled analogue from every other affinity molecule.

98. (new) The method of claim 40, wherein two or more conjugates are used, and wherein each affinity molecule in the two or more conjugates has a property capable of binding with the analyte in the sample and the labeled analyte at a different site on the analyte in the sample and a different site on the labeled analyte from every other affinity molecule, or each affinity molecule in the conjugate has a property capable of binding with the analyte in the sample and the labeled analogue at a different site on the analyte in the sample and a different

10/821,657 filed 04/08/2004

Wada et al.

Reply to Office Action of 07/24/2006

site on the labeled analogue from every other affinity molecule, and wherein the charged carrier molecule has a property capable of causing a change in a separation property of the labeled analyte or the labeled analogue by binding to the labeled analyte or the labeled analogue through the affinity molecule to form a complex of the labeled analyte or the labeled analogue, the affinity molecule, and the charged carrier molecule.

99. (new) The method of claim 41, wherein two or more affinity molecules are used, and wherein each affinity molecule has a property capable of binding with the analyte in the sample and the labeled analyte at a different site on the analyte in the sample and a different site on the labeled analyte from every other affinity molecule, or each affinity molecule has a property capable of binding with the analyte in the sample and the labeled analogue at a different site on the analyte in the sample and a different site on the labeled analogue from every other affinity molecule, and wherein the charged carrier molecule has a property capable of causing a change in a separation property of the labeled analyte or the labeled analogue by binding to the labeled analyte or the labeled analogue through the affinity molecule to form a complex of the labeled analyte or the labeled analogue, the affinity molecule, and the charged carrier molecule.

100. (new) The method of claim 42, wherein two or more affinity molecules are used, and wherein each affinity molecule has a property capable of binding with the analyte in the sample and the analyte bound to the charged carrier molecule at a different site on the analyte in the sample and a different site on the analyte bound to the charged carrier molecule from every other affinity molecule, or each affinity molecule has a property capable of binding with the analyte in the sample and the analogue bound to the charged carrier molecule at a different site on the analyte in the sample and a different site on the analogue bound to the charged carrier molecule from every other affinity molecule, and wherein the charged carrier molecule has a property capable of causing a change in a separation property of the first complex by binding to the analyte or the analogue to form a complex of the analyte, the affinity molecule, and the charged carrier molecule.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.